


Recommendations for Restraint Use in Patients with Traumatic Brain Injury



## References:

1. McNett, M., Sarver, W., & Wilczewski, P. (2012). The prevalence, treatment and outcomes of agitation among patients with brain injury admitted to acute care units. *Brain Injury*, 26(9), 1155-1162. <https://doi.org/10.3109/02699052.2012.667587>
2. Eskandari, F., Abdullah, K. L., Zainal, N. Z., & Wong, L. P. (2017). Use of physical restraint: Nurses' knowledge, attitude, intention and practice and influencing factors. *Journal of Clinical Nursing*, 26(23-24). 4479-4488. <https://doi.org/10.1111/jocn.13778>
3. Lombard, L. A., & Zafonte, R. D. (2005). Agitation after traumatic brain injury: Considerations and treatment options. *American Journal of Physical Medicine and Rehabilitation*, 84(10), 797-812. <https://doi.org/10.1097/01.pmh.0000179438.22235.08>
4. Centers for Disease Control and Prevention. (2022, July 18). Traumatic Brain Injury and Concussion. <https://www.cdc.gov/traumaticbraininjury>
5. Votruba, K. L., Rapport, L. J., Vangel, S. J., Hanks, R. A., Lequerica, A., Whitman, R. D., & Langenecker, S. (2008). Impulsivity and traumatic brain injury: The relations among behavioral observation, performance measures, and rating scales. *Journal of Head Trauma Rehabilitation*, 23(2), 65-73. <https://doi.org/10.1097/01.HTR.0000314525.93381.69>
6. Jacq, G., Melot, K., Bezou, M., Foucault, L., Courau-Courtois, J., Cavelot, S., Lang, A., Bedos, J. P., Le-Boeuf, D., Boussard, J. M., & Legriel, S. (2018). Music for pain relief during bed bathing of mechanically ventilated patients: A pilot study. *PLOS One*, 13(11), <https://doi.org/10.1371/journal.pone.0207174>.

### **Updated June 2024**

*Brandie Crawford, APRN, MSN, AGACNP-BC*

*Bethany Evans, RN, MSN, ACNP-BC*

*Sandra Tsay, MSN, RN, AGCNS-BC, CNE*

*Diana Hayes, ACNP-BC*

*Allison Chandler, ACNP-BP*

*Jennifer Beavers, PharmD, BCPS*

### **Revised:**

*June 2024*